

Benefits of a grape-rich diet on human health: a nutrigenomics study underlining the potential role of non-coding RNAs

Rosa Anna Milella¹, Marica Gasparro¹, Fiammetta Alagna^{1,2}, Maria Francesca Cardone¹, Silvia Rotunno^{1,3}, Concetta Tiziana Ammollo⁴, Fabrizio Semeraro⁴, Apollonia Tullo⁵, Flaviana Marzano⁵, Domenico Catalano⁶, Mario Colucci⁴, Domenica D'Elia⁶✉

¹Research Centre for Viticulture and Enology, Council for Agricultural Research and Economics, Turi, Bari, Italy

²ENEA Italian National Agency for New Technologies Energy and Sustainable Economic Development, Trisaia Research Center, Rotondella, Matera, Italy

³Institute for Sustainable Plant Protection, National Research Council, Torino, Italy

⁴Department of Biomedical Sciences and Human Oncology, University of Bari "Aldo Moro", Bari, Italy

⁵Institute of Biomembranes, Bioenergetics and Molecular Biotechnologies, National Research Council, Bari

⁶Institute for Biomedical Technologies, National Research Council, Bari, Italy

Competing interests: RAM none; MG none; FA none; MFC none; SR none; CTA none; FS none; AT none; FM none; DC none; MC none; DD none

Grape (*Vitis vinifera* L.) is one of the most typical fruit of the Mediterranean diet, characterised by high polyphenols content with marked antioxidant and anti-inflammatory activities (Goszcz *et al.*, 2017). By using transcriptional profiling techniques, it has become increasingly clear that polyphenols can influence the expression of genes. Many of these genes are key elements of cell signalling cascades (Spencer 2009; Fraga and Oteiza 2011) and regulatory non-coding RNAs (ncRNAs) (Budisan *et al.*, 2017).

A previous study showed that table grape extracts exert a marked antithrombotic activity *in vitro* (Ammollo *et al.*, 2017). To shed light on the molecular basis of grape intake effects on human health and investigate the potential role of ncRNAs, we carried out a nutrigenomics study (Milella *et al.*, 2020a and 2020b). To this aim, 20 healthy subjects were enrolled to follow a grape-rich diet for 21 days. The gene expression profiles of peripheral blood mononuclear cells (PBMCs) extracted from six of these subjects, after 21 days of fresh table grape-rich diet and after an additional 28-day washout, were analysed. The results showed 930 genes differentially expressed. Among these genes, more than 200 are long ncRNAs (lncRNAs), almost all downregulated after the washout period when polyphenols' direct effect is supposed to be completely exhausted.

The functional analysis of differentially expressed genes revealed significant changes in processes critical for organismal and cell wellbeing such as inflammation and immunity, thrombosis, DNA and protein repair, autophagy and mitochondrial biogenesis. From our analysis of lncRNAs down-regulated in our study, we observed that many of them are over-expressed in many types of tumours, chronic diseases due to the persistency of inflammation and metabolic syndromes such as obesity. Altogether, these findings provide exciting clues for the crucial role of ncRNAs in grape intake's long-term

effects on a series of biological processes. These lncRNAs would deserve to be further investigated for potential applications in the care of chronic diseases and cancer and their prevention.

Acknowledgements

This work was supported by a grant from MIUR (Ministero Italiano dell'Università e della Ricerca - project "PON02_00186_2937475 - PROALIFUN") and by the CNR (National Research Council) Flagship Project InterOmics.

References

1. Ammollo CT *et al.* (2017) Grape intake reduces thrombin generation and enhances plasma fibrinolysis. Potential role of circulating procoagulant microparticles. *J Nutr Biochem* **50**:66–73. <http://dx.doi.org/10.1016/j.jnutbio.2017.08.012>
2. Budisan L *et al.* (2017) Dietary Intervention by Phytochemicals and Their Role in Modulating Coding and Non-Coding Genes in Cancer. *Int J Mol Sci* **18**(6). <http://dx.doi.org/10.3390/ijms18061178>
3. Fraga CG, Oteiza PI (2011) Dietary flavonoids: Role of (-)-epicatechin and related procyanidins in cell signalling. *Free Radic Biol Med* **51**(4):813–23. <http://dx.doi.org/10.1016/j.freeradbiomed.2011.06.002>
4. Goszcz K, Duthie GG, Stewart D, Leslie SJ, Megson IL (2017) Bioactive polyphenols and cardiovascular disease: chemical antagonists, pharmacological agents or xenobiotics that drive an adaptive response? *Br J Pharmacol* **174**:1209–25. <http://dx.doi.org/10.1111/bph.13708>
5. Milella RA *et al.* (2020b) Gene expression signature induced by grape intake in healthy subjects reveals wide-spread beneficial effects on peripheral blood mononuclear cells. *Journal of Functional Food*, **64**:103705. <http://dx.doi.org/10.1016/j.jff.2019.103705>
6. Milella RA *et al.* (2020a) Microarray data and pathway analyses of peripheral blood mononuclear cells from healthy subjects after a three weeks grape-rich diet. *Data in Brief* **29**, 105278. <http://dx.doi.org/10.1016/j.dib.2020.105278>
7. Spencer JP (2009) Flavonoids and brain health: multiple effects underpinned by common mechanisms. *Genes Nutr*. **4**(4):243–50. <http://dx.doi.org/10.1007/s12263-009-0136-3>

© 2021 Milella *et al.*; the authors have retained copyright and granted the Journal right of first publication; the work has been simultaneously released under a Creative Commons Attribution Licence, which allows others to share the work, while acknowledging the original authorship and initial publication in this Journal. The full licence notice is available at <http://journal.embnet.org>.